




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Brain natriuretic peptide as a prognostic factor in COVID-19

Irving Elliot Ortiz y Cruz, Valeria Juárez García, Israel Nayensei Gil Velázquez, Pedro José Curi Curi

PII: S2603-9249(23)00023-X

DOI: <https://doi.org/10.1016/j.mcpsp.2023.100385>

Reference: MCP 100385

To appear in:

Received date: 3 April 2023

Accepted date: 3 May 2023

Please cite this article as: I.E. Ortiz y Cruz, V.J. García, I.N.G. Velázquez, et al., Brain natriuretic peptide as a prognostic factor in COVID-19, (2023), <https://doi.org/10.1016/j.mcpsp.2023.100385>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Brain natriuretic peptide as a prognostic factor in COVID-19

Péptido cerebral natriurético como factor pronóstico en COVID-19

Irving Elliot Ortiz y Cruz², Valeria Juárez García³, Israel Nayensei Gil Velázquez², Pedro José Curi Curi^{1*}

¹Dirección de Planeación, Enseñanza e Investigación, Hospital Regional de Alta Especialidad Ixtapaluca, Estado de México, México.

²Servicio de Medicina Interna, Hospital Regional de Alta Especialidad Ixtapaluca, Estado de México, México.

³Facultad de Estudios Superiores Zaragoza, Universidad Nacional Autónoma de México, México.

*Corresponding autor

E-mails:

Irving Elliot Ortiz y Cruz : drirvingortiz@gmail.com

Valeria Juárez García : wonejiithr@comunidad.unam.mx

Israel Nayensei Gil Velázquez: hillipg@icloud.com

Pedro José Curi Curi : pcunacuri001@gmail.com

Resumen

Introducción: Actualmente la infección por SARS-CoV-2 ha provocado la muerte de 6.5 millones de personas a nivel mundial. La COVID-19 es una enfermedad que afecta primordialmente al sistema respiratorio y puede llegar a provocar falla multiorgánica. Debido a la mortalidad que ha generado se han generado diferentes índices pronósticos para determinar que pacientes son más propensos a complicarse y fallecer. En BNP es una hormona peptídica sintetizada en los ventrículos del corazón y se ha usado como indicativo de insuficiencia cardiaca y como factor pronóstico en pacientes con choque séptico. Por lo tanto, se ha planteado su uso como factor pronóstico en pacientes que presentan COVID-19.

Materiales y métodos: Se llevó a cabo estudio retrospectivo de casos y controles que incluyó a 100 pacientes confirmados con infección por SARS-CoV-2 por PCR-RT, los cuales fueron egresados a domicilio o fallecido hasta el 30 de mayo de 2020. Utilizando el expediente clínico electrónico de estos pacientes, se obtuvieron datos demográficos, clínicos y bioquímicos para realizar la comparación entre los sobrevivientes y los pacientes que fallecieron. Se realizó análisis estadístico con curva ROC para determinar el nivel de BNP al ingreso, que se asoció con mayor mortalidad intrahospitalaria.

Resultados y conclusiones: De los 100 pacientes incluidos, 50 fueron egresados a domicilio y 50 fallecieron durante su estancia intrahospitalaria. El 87% de los pacientes presentaba al menos una comorbilidad, siendo la obesidad la más frecuente (38 pacientes, 38%), seguido de hipertensión (25 pacientes, 25%). Existió diferencia estadísticamente significativa entre ambos grupos en las siguientes características: edad, género masculino, saturación de oxígeno por pulsioximetría, conteo de leucocitos, recuento de neutrófilos, deshidrogenasa láctica y proteína C reactiva. Respecto al BNP, se encontró que un punto de corte mayor a 32 pg/ml puede utilizarse como factor predictor de mortalidad intrahospitalaria (AUC 0.751) con una sensibilidad de 60% y especificidad de 90%; así como odds ratio de 13.5 (IC 95%; 4.6-39.9).

Un nivel de BNP, al ingreso, mayor de 32 pg/ml se asocia con mayor mortalidad intrahospitalaria en pacientes con COVID-19.

Palabras clave: *Péptido natriurético cerebral, COVID-19* / mortalidad, Pronóstico*

Abstract

Introduction: COVID-19 is a disease that affects the respiratory tract and can potentially lead to multiple organ failure and death. Due to its high mortality, different prognostic indexes have been proposed in order to identify which patients are more prone to develop complications and death. We aimed to determine if BNP, which is a peptide hormone used as an indicator of heart failure and as a prognostic factor in patients with septic shock, may be used as well as a prognostic factor in patients with COVID-19.

Materials and methods: A retrospective case-control study was carried out at a single medical center, including 100 patients with SARS-CoV-2 infection diagnosed by PCR-RT. From the electronic clinical records, data was obtained in order to compare survivors with the died patients in terms of blood BNP levels.

Results: Of the 100 patients studied, 50 were discharged home and 50 died during their hospital stay. There was a statistically significant difference between both groups regarding age, male gender, oxygen saturation, leukocyte and neutrophil count, lactic dehydrogenase, and C-reactive protein. Regarding BNP, it was found that a cut-off point > 32 pg/ml can be used as a predictor of in-hospital mortality (AUC 0.751) with a sensibility of 60% and sensitivity of 90%; as well as an odds ratio of 13.5 (95% CI; 4.6-39.9).

Discussion: An admission BNP level > 32 pg/ml is associated with a higher in-hospital mortality in patients with COVID-19 and can be considered as a prognostic factor for this disease.

Key words: *Brain natriuretic peptide, COVID-19* / mortality, Prognosis*

Abreviaturas

BNP: Péptido natriurético cerebral

COVID-19: Enfermedad por coronavirus 2019

NT-proBNP: prohormona N-terminal del péptido natriurético cerebral

SaO₂: saturación de oxígeno

SARS-CoV-2: Coronavirus tipo 2

Abbreviations

BNP: Brain natriuretic peptide

COVID-19: Coronavirus Disease 2019

NT-proBNP: N-terminal prohormone of brain natriuretic peptide

SaO₂: Oxygen saturation

SARS-CoV-2: Coronavirus type 2

Introduction

Currently, coronavirus type 2 (SARS-CoV-2) has been implicated as the etiological agent of Coronavirus Disease 2019 (COVID-19). The infectious clinical spectrum of SARS-CoV-2 goes from an asymptomatic process to a severe respiratory failure that can lead to death. Effects of SARS-CoV-2 virus is due to its property to bind with the human angiotensin I converting enzyme 2 (ACE2) receptor. Infections with this virus can lead to multiple organ failure because ACE2 receptor is expressed in almost all organs of the human body (1).

Brain natriuretic peptide (BNP) is a hormone mainly produced at the heart ventricles (2). Its synthesis and release is secondary to heart wall distension, ventricular dilation and/or increased pressure in the cardiac chambers. BNP elevated blood levels promote vasodilation, increased diuresis, and produces suppression of the renin-angiotensin-aldosterone system in order to decrease heart preload (3).

BNP, which is a recognized heart failure biomarker, has also been used as a prognostic factor in patients with sepsis and septic shock, regardless of their previous cardiovascular condition. This study aims to determine if BNP can be also used as a prognostic factor in patients with SARS-CoV-2 infection.

Methodology

A retrospective, observational, analytical, and cross-sectional case-control clinical study was designed, in which all patients over 18 years with a diagnosis of COVID-19 confirmed by reverse transcriptase polymerase chain reaction (PCR-RT) were included. Patients were admitted to the internal medicine service of a third level hospital in a one month period during the 2020 pandemic. Those with a previous diagnosis of chronic heart failure, chronic kidney disease, hyperthyroidism, chronic liver disease, malignancy, and a history of subarachnoid hemorrhage in the last 3 months were excluded. Chronic kidney disease during hospital stay, serum creatinine levels >1.5 mg/dl or blood glucose >250 mg/dl were considered as elimination criteria for this study. Patients were divided in two groups: the ones who were discharged home due to clinical improvement (controls), and those who died during their hospital stay (cases). Clinical background records were collected, specially those referred to comorbidities and

medication intake, as well as vital signs and anthropometric measurements to determine body mass index (BMI). Laboratory studies such as hematic biometrics, blood chemistry, liver function tests, ultrasensitive C-reactive protein, procalcitonin, ferritin, and blood BNP levels were considered.

Clinical and laboratory data were collected from clinical records within the first 24 hours of hospital admission. This information was recorded into an Excel spreadsheet for further analysis using SPSS version 21 software. Quantitative variables are expressed as mean \pm standard deviation and categorical variables are expressed as frequency and percentage in relation to the population at risk. BNP values were analyzed using the ROC curve and then Odds ratio (OR) values were calculated.

Results

A sample of 100 patients was obtained out of total of 204 clinical records studied. Fifty patients were randomly assigned to each study group (cases and controls) using an Urna Software.

Table 1 shows sociodemographic variables, somatometry and vital signs. Male gender was predominately observed at the problem group, which were eight years older in relation to the control group. Despite the fact that there were no significant differences in comorbidities and somatometry, the most frequent comorbidities in both groups were obesity, arterial hypertension and type 2 diabetes mellitus. There were no differences in vital signs, diastolic blood pressure, heart rate, respiratory rate and temperature at both studied groups. However, systolic blood pressure was higher in the problem group and oxygen saturation (SpO_2) was higher in the control group. In **table 2** laboratory values are shown, and we can see higher levels in the problem group related to hemoglobin, leukocytes and absolute neutrophils, as well as blood urea nitrogen (BUN), levels of lactic dehydrogenase (LDH) and C-reactive protein.

Figure 1 shows a ROC curve for BNP values of the problem group. It was found that the cut-off point for serum BNP was 32 pg/ml, which represented the highest sensibility and specificity for the end point of death in patients with COVID-19. Therefore, a patient who died showed a 60% probability of having a serum BNP level > 32 pg/mL at admission. A patient who was discharged from hospital showed a 90% probability of

having an initial BNP serum level < 32 pg/ml. It was determined that the positive predictive value of BNP for intrahospital mortality from COVID-19 is 0.85 and the negative predictive value is 0.69. The area under the curve (AUC) of the BNP as a prognostic tool for in-hospital mortality due to COVID-19 was 0.751. Since the confidence interval does not include the value of 0.50, we can affirm that the AUC of the BNP concentration is different from non-discrimination.

Table 3 shows the number of patients in both groups who presented both higher and lower BNP values on admission, taking 32 pg/ml as the cut-off point. The calculated OR was 13.5, so that patients with COVID-19 and $\text{BNP} \geq 32$ pg/ml are 13 times more likely to die during their hospital stay compared to those patients with a BNP level below this cut-off point. In turn, the 95% confidence interval was between 4.6 and 39.9.

Figure 2 shows that most of the patients in the problem group had BNP concentrations ≥ 32 pg/ml compared to the control group. Positive and negative odds ratios were calculated. The positive likelihood ratio (CPR) resulted in 6, while the negative likelihood ratio (CPN) was obtained in 0.44. **Figure 3** shows the Fagan nomogram. The first line shows the pre-test probability, which is represented by intra-hospital mortality in patients with COVID-19. Starting from the prevalence value, which in this study is 50, a line was drawn that intersects the values of the probability ratios, both positive and negative, and finally extends to the post-test probability axis. A positive post-test probability value of 86% was obtained. The negative post-test probability value was 31%. Therefore, in a clinical scenario with the prevalence of our study, a patient with COVID-19 and BNP on admission greater than 32 pg/ml has an 86% probability of dying and, conversely, if the BNP result is less than this cut-off value, the patient has a 31% probability of being discharged at home.

Discussion

Morbidity and mortality due to COVID-19 disease increased significantly all worldwide. Therefore, it was necessary to develop scales and search for prognostic factors that would allow the identification of patients with the highest risk for complications and death. BNP is a biomarker that has been used as a prognostic factor in patients with sepsis and septic shock, because its short term elevation is a mortality predictor (4).

Currently, there are few researches related to the prognostic value of BNP as a biomarker in COVID-19, since the N-terminal prohormone of brain natriuretic peptide (NT-proBNP) has mostly been used as a prognostic biomarker. Elevated NT-proBNP concentrations at admission in COVID-19 patients have been shown to be associated with higher rates of mechanical ventilation, intensive care unit (ICU) admissions, and a twice probability risk of in-hospital mortality, regardless of a clinical background of a previous heart failure (5)(6). It has been stated that the NT-proBNP cut-off point to predict mortality in patients with COVID-19 is 88.64 pg/mL, with a 100% sensibility and a 66.67% sensitivity. Patients with NT-proBNP blood concentrations greater than that value have a lower cumulative survival rate compared to those with lower levels (7). In the present study, BNP was used as a mortality predictor biomarker that showed a relationship between elevated levels and in-hospital mortality in patients with COVID-19, with a 32 pg/ml as a cut-off point. The elevation of BNP/NT-pro-BNP has not yet been studied, but it is thought to be multifactorial (8). Currently, viral RNA has been associated with direct damage of the virus at the myocardial tissue. This tropism is associated with the expression of the angiotensin-converting enzyme receptor 2 (ACE2) in cardiac cells, which allows SARS-CoV-2 to infect and replicate within cardiomyocytes, leading to infiltration of monocytes, lymphocytes and plasma cells, and developing myocarditis due to COVID-19 with increased cardiac biomarkers (9) (10).

In addition to the direct cytopathic effect of SARS-CoV-2 on cardiomyocytes, it has also been proposed that myocardial damage is mediated by the uncontrolled release of cytokines. Cytokine storm is the main cause of severity and death in patients with COVID-19. It is the consequence of an excessive immune response that can be fatal. There is an imbalance in the cellular response of T helper 1 lymphocytes, excessive production of proinflammatory cytokines (Interleukin-6, tumor necrosis factor- α) and chemical mediators, that are capable of damaging the myocardium and causing cardiac dysfunction (11) (10). In the problem group, the numbers of leukocytes, absolute neutrophils, and ultrasensitive C-reactive protein levels were significantly higher, which is associated with a great inflammatory response induced by SARS-CoV-2 infection. These data also show that myocardial injury and the consequent elevation of B-type natriuretic peptide are strongly associated with an inflammatory pathogenic substrate.

Leukocytes are attracted by inflammatory signals to cardiac tissue, producing numerous reactive oxygen species at the site of injury and increased oxidative stress, which leads to cardiomyocyte deterioration associated with modification of contractile proteins and dysregulation of cellular redox states. These changes promote apoptosis of cardiomyocytes and determine a state of reparative fibrosis (12)(13).

In this study, no statistically significant difference was documented in lymphocyte count between both groups. Our data differ from most published studies in which lymphopenia is a typical profile in patients with COVID-19 (14). Lymphopenia represents an independent risk factor for mortality and has even become part of some indices or scores that try to predict the progression of COVID-19 towards severe forms, such as the CALL score or the COVID-gram score (15)(16).

Multiple risk factors leading to severe COVID-19 disease have currently been identified. Advanced age (> 65 years), male gender, chronic degenerative diseases (diabetes, arterial hypertension, obesity, and cardiovascular diseases) are significant risk factors for disease severity, complications, and poor prognosis. Since the beginning of the pandemic, it has been determined that older people have a more severe presentation of the disease from COVID-19 and a higher mortality. The processes of aging and cellular senescence of the immune system are factors that leave the elderly patient in a vulnerable state (17). In this study it was observed that in the problem group most of the patients were in the fifth decade of life and although they were not older than 60 years, it is corroborated that age is a prognostic factor for the outcome of COVID-19. Age has a directly proportional relationship with the increase in mortality. This is evident in patients ≥ 60 years, where the highest mortality occurs in patients ≥ 80 years in whom the risk of death is six times higher compared to young patients (18). Similarly, it has been shown that the male gender predicts a higher mortality rate compared to the female gender (19). In the problem group, the number of men was significantly higher compared to the control group, which supports that the male gender is a predictor of increased risk of death in adults with COVID-19. Sex hormones are involved in the immune response to SARS-CoV-2 infection. In women, estrogens are a protective factor and in infectious disease promote the proliferation of T cells and therefore a stronger immune response. In men, androgens, such as testosterone and

dihydrotestosterone, increase the count and function of the main cells responsible for cytokine storm syndrome, the neutrophils. Therefore, there is a greater predisposition to develop an exaggerated immune response and greater complications. In general, male sex hormones facilitate the entry of the virus into tissues, since they increase the activity of the ACE2 receptor and favor the expression of the transmembrane protease serine 2 (TMPRSS2) (20)(21).

Regarding vital signs, the problem group presented significantly lower SaO_2 . Hypoxemia occurs when the oxygen content in arterial blood is decreased. Hypoxemia in patients with COVID-19 is a consequence of alterations between ventilation/perfusion (V/Q) caused by the presence of pulmonary edema and loss of alveolar elasticity, which produces alveolar collapse and therefore alterations in hematosis (22). In patients with COVID-19, the severity of hypoxemia is an independent predictor of mortality, respiratory complications, and use of mechanical ventilation. In the presence of hypoxia, various compensatory mechanisms occur, such as the increase in the red blood cell count. Under hypoxic conditions, the juxtaglomerular cells of the kidney are stimulated by factor 1- α to favor the secretion of erythropoietin, which in the bone marrow favors the production of red blood cells, of which main component is hemoglobin (23). This explains why the patients in the problem group with considerably low oxygen saturation have significantly higher hemoglobin levels compared to the control group.

This study becomes important as no articles have been published that relate B-type natriuretic peptide with in-hospital mortality in COVID-19 population. It suggests, as well, a new low-cost and usefull prognostic factor for SARS-CoV-2 infection.

Tables and figures

Variables	CONTROL group n (%) / Mean \pm SD (Min - Max)	PROBLEM group n (%) / Mean \pm SD (Min - Max)	p
Age (years)	45 \pm 12 (19-64)	53 \pm 9 (26 - 65)	0.0003
Gender			
Male	24 (24%)	35 (35%)	0.042
Feminine	26 (26%)	15 (15%)	
Comorbidity			
SAH	11 (11%)	14 (14%)	NS

DM2	10 (10%)	14 (14%)	NS
Obesity	21 (21%)	17 (17%)	NS
Smoking	3 (3%)	4 (4%)	NS
Other	4 (4%)	4 (4%)	NS
Somatometry			
Weight (kg)	81 ± 23 (56-129)	84 ± 19 (54-150)	NS
Height (m)	1.62 ± 0.33 (1.46-1.80)	1.63 ± 0.10 (1.42-1.75)	NS
BMI	31 ± 8 (20-50)	32 ± 16 (22-55)	NS
Vital signs			
Systolic BP	114 ± 16 (90-160)	122 ± 16 (90 - 157)	0.0141
Diastolic BP	72 ± 11 (56-100)	75 ± 10 (60 - 100)	NS
HR			
(beats/min)	107 ± 20 (63 - 189)	105 ± 17 (62 - 157)	NS
RR			
(breaths/min)	27 ± 7 (18 - 48)	29 ± 6 (21 - 48)	NS
Temperature			
(°C)	37.5 ± 1 (35.6 - 41)	37.2 ± 0.8 (35.6 - 38.8)	NS
O ₂ saturation			
(%)	79 ± 16 (36 - 98)	66 ± 16 (21 - 91)	0.0001

Table 1. Comparison of the sociodemographic variables, somatometry and vital signs of the studied groups

Abbreviations: SAH=systemic arterial hypertension, DM2=type 2 diabetes mellitus, Kg=kilograms, m=meters, BMI=body mass index, BP=blood pressure, HR=heart rate, min=minutes, RR= respiratory rate, °C=Celsius degrees and O₂=oxygen

Variables	CONTROL group Mean \pm SD (Min - Max)	PROBLEM group Mean \pm SD (Min - Max)	p
Blood count			
Hemoglobin (g/dl)	14.3 \pm 2 (8.7 - 17.7)	15.1 \pm 1.8 (8.1 - 18.5)	0.0381
Hematocrit (%)	42.4 \pm 6.2 (25.2 - 57.5)	44.4 \pm 5.2 (25.4 - 56)	NS
MCV	88.5 \pm 3.7 (77.3 - 98.9)	89.4 \pm 5 (73.3 - 100.4)	NS
MCH	30 \pm 1.7 (23.8 - 33.9)	30.3 \pm 1.9 (23.8 - 34.2)	NS
Leu (cell/mm ³)	8,838 \pm 3,795 (14 - 19,800)	11,648 \pm 5,150 (4,200 - 27,600)	0.0025
Neu (cell/mm ³)	7,366 \pm 3,507 (6 - 17,030)	10,134 \pm 4,826 (2,410 - 25,780)	0.0014
Lymph (cell/mm ³)	1,158 \pm 1,331 (170 - 9,680)	906 \pm 92 (90 - 2,430)	NS
Plaq (cell/mm ³)	270,300 \pm 125,597 (48,000 - 812,000)	265,240 \pm 112,523 (60,000 - 677,000)	NS
Blood chemistry			
Glucose (mg/dl)	116 \pm 51 (55 - 250)	164 \pm 95 (73 - 250)	NS
BUN (mg/dl)	13.9 \pm 6 (5.7 - 33.3)	18.9 \pm 8 (6.4 - 45.3)	0.0006
Creatinine (mg/dl)	0.8 \pm 0.2 (0.4 - 1.1)	0.9 \pm 0.3 (0.5 - 1.4)	NS
Liver function tests			
Alb (g/dl)	3.5 \pm 0.5 (2.2 - 4.3)	3.5 \pm 0.4 (2.4 - 4.4)	NS
GOT (IU/L)	43 \pm 22 (8 - 115)	63 \pm 111 (16 - 787)	NS
TGP (IU/L)	42 \pm 26 (3 - 141)	51 \pm 53 (12 - 358)	NS
ALP (mg/dl)	109 \pm 53 (53 - 228)	129 \pm 81 (58 - 454)	NS
GGT (IU/L)	145 \pm 164 (12 - 738)	134 \pm 133 (28 - 711)	NS
DHL (IU/L)	384 \pm 103 (173 - 597)	489 \pm 244 (208 - 1,794)	0.0061
Proca (ng/ml)	0.63 \pm 1.14 (0.05 - 7.35)	0.93 \pm 1.48 (0.06 - 7.05)	NS
MGV (mm/h)	30 \pm 17 (1 - 61)	31 \pm 15 (1 - 65)	NS
PCR (mg/l)	175.1 \pm 106.19 (5.07 - 437.4)	237.5 \pm 102.74 (34.95 - 478.46)	0.0036
Ferritin (ng/ml)	854.2 \pm 856.78 (41.39 - 3,429.84)	2,051.85 \pm 5,411.35 (60.94 - 38,746)	NS

Table 2. Laboratory variables of the study groups.

Abbreviations: MCV=mean corpuscular volume, MCH=mean corpuscular hemoglobin; Leu=leukocytes, Neu=absolute neutrophils, Lymph=absolute lymphocytes, Plaq=platelets, BUN=blood urea nitrogen, Alb=albumin, GOT=glutamic-oxaloacetic transaminase, GGT=glutamic-pyruvic transaminase, ALP=alkaline phosphatase, GGT= gamma- glutamyl transpeptidase, LDH=lactic dehydrogenase, Proca=procalcitonin, ESR=erythrocyte sedimentation rate, CRP= c-reactive protein, g=grams, dl=deciliter, mm=millimeters, mg=milligrams, IU= international units, L=liters, ng=nanograms, h=hour

BNP	Problem	Control	p	OR (95% CI)
≥ 32 pg/ml	30 (30%)	5 (5%)	< 0.0001	13.5 (4.6-39.9)
< 32 pg/ml	20 (20%)	45 (45%)		

Table 3. Contingency table of the frequency of BNP values at the cut-off point established by the ROC curve in both study groups.

Abbreviations: pg=picograms, ml=milliliters, OR=odds ratio, CI=confidence interval




Figure 1. ROC curve for BNP values in the group of deceased patients
 Abbreviations: ROC=Receiver Operating Characteristic, AUC=area under the curve, CI=confidence Interval




Figure 2. Grouped bar chart and its comparison with the BNP values at the cut-off point established by the ROC curve.





Figure 3. Fagan nomogram summarizing the characteristics of BNP as a prognostic study of in-hospital mortality in patients with COVID-19

References

1. Devaux CA, Rolain JM, Raoult D. ACE2 receptor polymorphism: Susceptibility to SARS-CoV-2, hypertension, multi-organ failure, and COVID-19 disease outcome. Vol. 53, *Journal of Microbiology, Immunology and Infection*. Elsevier Ltd; 2020. p. 425–35.
2. Shen S, Ye J, Wu X, Li X. Association of N-terminal pro-brain natriuretic peptide level with adverse outcomes in patients with acute myocardial infarction: A meta-analysis. Vol. 50, *Heart and Lung*. Mosby Inc.; 2021. p. 863–9.
3. Nakagawa Y, Nishikimi T, Kuwahara K. Atrial and brain natriuretic peptides: Hormones secreted from the heart. Vol. 111, *Peptides*. Elsevier Inc.; 2019. p. 18–25.
4. Vallabhajosyula S, Wang Z, Murad MH, Vallabhajosyula S, Sundaragiri PR, Kashani K, et al. Natriuretic Peptides to Predict Short-Term Mortality in Patients With Sepsis: A Systematic Review and Meta-analysis. *Mayo Clin Proc Innov Qual Outcomes*. 2020 Feb;4(1):50–64.
5. O'donnell C, Ashland MD, Vasti EC, Lu Y, Chang AY, Wang P, et al. N-Terminal Pro-B-Type Natriuretic Peptide as a Biomarker for the Severity and Outcomes With COVID-19 in a Nationwide Hospitalized Cohort. *Journal of the American Heart Association J Am Heart Assoc* [Internet]. 2021;10:22913. Available from: <https://www.ahajournals.org/>
6. Pranata R, Huang I, Lukito AA, Pahajjo SB. Elevated N-terminal pro-brain natriuretic peptide is associated with increased mortality in patients with COVID-19: systematic review and meta-analysis. *Postgrad Med J* [Internet]. 2020;96:387–91. Available from: <http://pmj.bmj.com/>
7. Gao L, Jiang D, Wen X song, Cheng X cheng, Sun M, He B, et al. Prognostic value of NT-proBNP in patients with severe COVID-19. Available from: <https://doi.org/10.1186/s12931-020-01352-w>
8. Junior GLG de A, Braga F, Jorge JK, Nobre GF, Kalichsztein M, de Faria P de MP, et al. Prognostic value of troponin-t and b-type natriuretic peptide in patients hospitalized for covid-19. *Arq Bras Cardiol*. 2020 Oct 1;115(4):660–6.
9. Bailey AL, Dmytrenko O, Greenberg L, Bredemeyer AL, Ma P, Liu J, et al. SARS-CoV-2 Infects Human Engineered Heart Tissues and Models COVID-19 Myocarditis. *JACC Basic Transl Sci*. 2021 Apr 1;6(4):331–45.
10. Tavazzi G, Pellegrini C, Maurelli M, Belliato M, Sciutti F, Bottazzi A, et al. Myocardial localization of coronavirus in COVID-19 cardiogenic shock. *Eur J Heart Fail* [Internet]. 2020;22:911–5. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/ehf.1828>
11. Hu B, Huang S, Yin | Lianghong. The cytokine storm and COVID-19. *J Med Virol* [Internet]. 2021;93:250–6. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/jmv.26232>
12. Tangos M, Budde H, Kolijn D, Sieme M, Zhazykbayeva S, Lódi M, et al. SARS-CoV-2 infects human cardiomyocytes promoted by inflammation and oxidative stress. *Int J Cardiol*. 2022 Sep 1;362:196–205.

13. Siripanthong B, Nazarian S, Muser D, Deo R, Santangeli P, Khanji MY, et al. Recognizing COVID-19–related myocarditis: The possible pathophysiology and proposed guideline for diagnosis and management. *Heart Rhythm*. 2020 Sep 1;17(9):1463–71.
14. Xiao Hua Luo. Yan Zhu. Jian Mao. Rui Chan Du. T cell immunobiology and cytokine storm of COVID-19. *Scand J Immunol*. 2020;
15. Ji D, Zhang D, Xu J, Chen Z, Yang T, Zhao P, et al. Prediction for Progression Risk in Patients With COVID-19 Pneumonia: The CALL Score. *Clin Infect Dis* [Internet]. 2020; Available from: <https://academic.oup.com/cid/article/71/6/1393/5818317>
16. Liang W, Liang H, Ou L, Chen ; Binfeng, Chen A, Li C, et al. Development and Validation of a Clinical Risk Score to Predict the Occurrence of Critical Illness in Hospitalized Patients With COVID-19. 2020; Available from: <http://118.126.104.170/>
17. Wirth R, Becker · C, Djukic · M, Drebenstedt · C, Hepner · H J, Jacobs · A H, et al. Gerontologie+Geriatric Übersichten COVID-19 im Alter-Die geriatrische Perspektive Einleitung. *Z Gerontol Geriatr* [Internet]. 2021;14:152–60. Available from: <https://doi.org/10.1007/s00391-021-01864-0>
18. Bonanad C, García-Blas S, Tarazona-Santabalbina F, Sanchis J, Bertomeu-González V, Fácila L, et al. The Effect of Age on Mortality in Patients With COVID-19: A Meta-Analysis With 611,583 Subjects. *J Am Med Dir Assoc*. 2020 Jul 1;21(7):915–8.
19. Nguyen NT, Chinnid J, de Ferrante M, Kiraly KA, Hohmann SF, Amin A. Male gender is a predictor of higher mortality in hospitalized adults with COVID-19. 2021; Available from: <https://doi.org/10.1371/journal.pone.0234066>
20. Moradi F, Enjezab B, Ghadiri Azad A. The role of androgens in COVID-19. Vol. 14, *Diabetes and Metabolic Syndrome. Clinical Research and Reviews*. Elsevier Ltd; 2020. p. 2003–6.
21. Wambier CG, Goren A, Vaño-Galván S, Paulo J, Ramos M, Ossimetha A, et al. Androgen sensitivity gateway to COVID-19 disease severity. 2020; Available from: <https://onlinelibrary.wiley.com/doi/10.1002/ddr.21688>
22. Dhont S, Derom F, van Braeckel E, Depuydt P, Lambrecht BN. Conceptions of the pathophysiology of happy hypoxemia in COVID-19. Vol. 22, *Respiratory Research*. 2021.

Central illustration



Conflicto de intereses

Los autores declaran no tener ningún conflicto de intereses en relación con este artículo.